



Determinant majorization and the work of Guo-Phong-Tong and Abja-OLive

F. Reese Harvey¹ · H. Blaine Lawson Jr.²

Received: 16 September 2022 / Accepted: 31 March 2023

© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2023

Abstract

The objective of this note is to establish the Determinant Majorization Formula $F(A)^{\frac{1}{N}} \geq \det(A)^{\frac{1}{n}}$ for all operators F determined by an invariant Gårding-Dirichlet polynomial of degree N on symmetric $n \times n$ matrices. Here *invariant* means under the group $O(n)$, $U(n)$ or $Sp(n) \cdot Sp(1)$ when the matrices are real symmetric, Hermitian symmetric, or quaternionic Hermitian symmetric respectively. We also establish this formula for the Lagrangian Monge Ampère Operator. This greatly expands the applicability of the recent work of Guo-Phong-Tong and Guo-Phong for differential equations on complex manifolds. It also relates to the work of Abja-Olive on interior regularity. Further applications to diagonal operators and to operators depending on the ordered eigenvalues are given. Examples showing the precision of the results are presented. For the application to Abja-Olive's work, and other comments in the paper, we establish some results for Gårding-Dirichlet operators in appendices. One is an exhaustion lemma for the Gårding cone. Another gives bounds for higher order derivatives, which result from their elegant expressions as functions of the Gårding eigenvalues. There is also a discussion of the crucial assumption of the Central Ray Hypothesis.

Contents

1	Introduction	...
2	The Basic Lemma-majorization of the determinant	...
3	Gårding-Dirichlet polynomial operators	...
4	Invariant G-D operators	...
5	Relevance to the work of Guo-Phong-Tong and Guo-Phong	...
6	Interior regularity-work of Abja-Olive	...

Communicated by Sun-Yung Alice Chang, Ph.D.

Second author was partially supported by the Simons Foundation.

H. Blaine Lawson Jr.
blaine@math.sunysb.edu

F. Reese Harvey
harvey@rice.edu

¹ Rice University, Houston, TX, USA

² Stony Brook University, Stony Brook, NY, USA

7	On the enormous universe of invariant G-D operators
8	Determinant majorization for the Lagrangian Monge-Ampère operator
9	Being a Gårding-Dirichlet operator is not enough for determinant majorization
10	Other applications of the Basic Lemma
Appendix A: The interior of the polar cone	
Appendix B: The Gårding gradient map	
Appendix C: The exhaustion Lemma	
Appendix D: The Güler derivative estimates for G-D operators	
Further discussion—second derivative formulas	
Appendix E: The central ray hypothesis	
References	

1 Introduction

In a recent fundamental paper [10], B. Guo, D. H. Phong, and F. Tong established an *apriori* L^∞ estimate for the complex Monge-Ampère equation on Kähler manifolds by purely PDE techniques. Their main theorem applied to many other equations as well. The basic hypothesis in the theorem can be established by proving a determinantal majorization inequality for the operator. In a later paper Guo and Phong proved many further results on Hermitian manifolds with the same hypothesis [11] (see also [12]). The point of this paper is to *prove the determinantal majorization inequality for every invariant Gårding-Dirichlet operator*, which means that all the results above apply to this very large family of differential equations. This inequality also appears as a hypothesis in work of S. Abja, S. Dinew and G. Olive [1, 2] on regularity, where there is also a compactness hypothesis which we establish in some generality. So their results also apply to this same large constellation of equations.

The operators in question can be defined as follows. Let $p(\lambda_1, \dots, \lambda_n)$ be a real homogeneous polynomial of degree N on \mathbb{R}^n which is symmetric in the variables $\lambda_1, \dots, \lambda_n$. We assume that all the coefficients of p are ≥ 0 .

We now define a polynomial operator $F : \text{Sym}^2(\mathbb{R}^n) \rightarrow \mathbb{R}$, of degree N on symmetric $n \times n$ -matrices, by

$$F(A) \equiv p(\lambda_1(A), \dots, \lambda_n(A)) \tag{1.1}$$

where $\lambda_1(A), \dots, \lambda_n(A)$ are the eigenvalues of $A \in \text{Sym}^2(\mathbb{R}^n)$. Note that $p(\lambda_1(A), \dots, \lambda_n(A))$ only makes sense because the polynomial $p(\lambda_1, \dots, \lambda_n)$ is symmetric, i.e., invariant under permutations of the λ_j 's. Since $\lambda(A) = \{\lambda_1(A), \dots, \lambda_n(A)\}$ is invariant under the action of $O(n)$ acting by conjugation on $\text{Sym}^2(\mathbb{R}^n)$, $F(A)$ is also $O(n)$ -invariant. This means that F defines an operator on every riemannian manifold by using the riemannian hessian [15].

We can do the analogous thing in the complex case. Let $A_{\mathbb{C}}$ be a hermitian symmetric $n \times n$ -matrix with eigenvalues $\lambda_k(A_{\mathbb{C}})$ for $k = 1, \dots, n$. Taking p as in the above real case, we can define

$$F(A_{\mathbb{C}}) \equiv p(\lambda_1(A_{\mathbb{C}}), \dots, \lambda_n(A_{\mathbb{C}})). \tag{1.2}$$

This operator is invariant under the unitary group $U(n)$ and makes sense on any Hermitian complex (or almost complex) manifold by using $i\partial\bar{\partial}u$ and the metric.

Similar remarks can be made in the quaternionic case.

In these cases we shall prove the determinantal majorization inequalities:

$$\begin{aligned} F^{\frac{1}{N}}(A) &\geq F(I)^{\frac{1}{N}} \det(A)^{\frac{1}{n}} \\ F^{\frac{1}{N}}(A_{\mathbb{C}}) &\geq F(I)^{\frac{1}{N}} \det_{\mathbb{C}}(A_{\mathbb{C}})^{\frac{1}{n}} \\ F^{\frac{1}{N}}(A_{\mathbb{H}}) &\geq F(I)^{\frac{1}{N}} \det_{\mathbb{H}}(A_{\mathbb{H}})^{\frac{1}{n}} \end{aligned} \quad (1.3)$$

where $F(I) > 0$. All three follow immediately from Basic Lemma 2.1.

We shall also prove this inequality for the Lagrangian Monge-Ampère operator M_{Lag} which is also defined on any almost complex Hermitian manifold (in particular, on symplectic manifolds with a Gromov metric). The defining polynomial is unitarily invariant, but the proof here is different because this polynomial is zero on all traceless Hermitian matrices, and so it is out of the category of operators considered in (1.3). In Sect. 8 we prove that in complex dimension n ,

$$M_{\text{Lag}}(A)^{\frac{1}{2n}} \geq \det_{\mathbb{R}}(A)^{\frac{1}{2n}}. \quad (1.4)$$

See [18] for complete details concerning this new operator.

The inequalities (1.3) also hold for polynomials p which are not symmetric but satisfy the Central Ray Hypothesis in Sect. 2. One just orders the eigenvalues $\lambda_1(A) \leq \dots \leq \lambda_n(A)$ and defines $F(A)$ by (1.1). Here F is not necessarily a polynomial. (See Sect. 10.)

Determinant majorization is also established for *diagonal operators*

$$p(a_{11}, a_{22}, \dots, a_{nn})$$

where p satisfies the hypotheses of Basic Lemma 2.1. However, if one drops the Central Ray Hypothesis, determinant majorization can fail. We show that for examples of this type local interior regularity can also fail. This is done in Sect. 9.

As mentioned before the family of polynomial operators F for which the determinant majorization holds is huge. One way to see this is to consider Gårding-Dirichlet (or G-D) operators on $\text{Sym}^2(\mathbb{R}^n)$, which are invariant. These are homogeneous real polynomials F on symmetric matrices with $F(I) > 0$ such that:

- (i) $t \mapsto F(tI + A)$ has all real roots for every A ,
- (ii) The Gårding cone Γ contains $\{A : A > 0\}$, where Γ is defined as the connected component of $\text{Sym}^2(\mathbb{R}^n) - \{F = 0\}$ containing the identity I ,
- (iii) F is invariant under the action of $O(n)$ by conjugation on $\text{Sym}^2(\mathbb{R}^n)$.

This family contains all elementary symmetric functions, the p -fold sum operator (Example 3.3), and many more (see [[16], §5]). The set of these operators is closed under products, under directional derivatives in Gårding cone directions, and under a certain “composition” rule (see Sect. 7).

Every such operator F can be expressed, in terms of the eigenvalues $\lambda(A)$ of A , as $F(A) = p(\lambda(A))$ where $p(\lambda_1, \dots, \lambda_n)$ is a symmetric homogeneous polynomial which is > 0 on $\mathbb{R}_{>0}^n$. After normalizing by a positive constant, p satisfies the three Assumptions of the Basic Lemma 2.1. Assumption (1) follows from this positivity, and Assumption (3) follows from the $O(n)$ -invariance. This polynomial p transfers over directly to the complex and quaternionic cases by applying it to the eigenvalues of $A_{\mathbb{C}}$ or $A_{\mathbb{H}}$, and so (1.3) holds.

We want to point out that all the operators considered in this paper are elliptic. In particular if F is a G-D polynomial operator with Gårding cone Γ , then a strengthened form of weak ellipticity, namely $\langle \nabla F, P \rangle > 0$ for all $P > 0$, holds in its domain Γ .

For the applications to the work of [1, 2] we need an Exhaustion Lemma which is proved in Appendix C. For the log of G-D polynomials there are elegant higher order estimates which we thought would be good to include here (Appendix D). An important assumption in the Basic Lemma is the Central Ray Hypothesis, which is discussed from several points of view in Appendix E. In a forthcoming article we shall give a comprehensive introduction to G-D operators and Gårding theory. However, in this paper the focus is centered on the following Basic Lemma 2.1.

We want to express our gratitude to Guillaume Olive for his careful reading of and helpful remarks on the first version of this paper.

2 The Basic Lemma–majorization of the determinant

Let p ($\not\equiv 0$) be a real homogeneous polynomial of degree N on \mathbb{R}^n , and set $e \equiv (1, 1, \dots, 1) \in \mathbb{R}^n$. We write

$$p(x) = \sum_{\alpha} a_{\alpha_1 \dots \alpha_n} x_1^{\alpha_1} \dots x_n^{\alpha_n} = \sum_{|\alpha|=N} a_{\alpha} x^{\alpha}.$$

Assumptions

- (1) All coefficients $a_{\alpha} \geq 0$.
- (2) (Normalization)

$$p(e) = \sum_{\alpha} a_{\alpha} = 1.$$

- (3) The crucial Central Ray Hypothesis:

$$D_e p = k e \quad \text{for } k > 0,$$

i.e.,

$$\frac{\partial p}{\partial x_1}(e) = \dots = \frac{\partial p}{\partial x_n}(e) = k > 0.$$

Note that

$$\frac{\partial p}{\partial x_j}(x) = \sum_{\alpha} a_{\alpha} \alpha_j x_1^{\alpha_1} \dots x_j^{\alpha_j-1} \dots x_n^{\alpha_n} \quad (2.1)$$

so that

$$\frac{\partial p}{\partial x_j}(e) = \sum_{\alpha} a_{\alpha} \alpha_j, \quad \text{for each } j = 1, \dots, n. \quad (2.2)$$

Therefore, (3) can be restated as

(3)' The Central Ray Hypothesis:

$$\sum_{|\alpha|=N} a_{\alpha} \alpha_j = k \quad \text{for each } j = 1, \dots, n \text{ with } k > 0.$$

It turns out that the following lemma goes back to a paper of Leonid Gurvits [13] in combinatorics.

BASIC LEMMA 2.1 For all $x_1 > 0, \dots, x_n > 0$,

$$p(x)^{\frac{1}{N}} \geq (x_1 \cdots x_n)^{\frac{1}{n}}.$$

Proof We first note that:

$$\begin{aligned} \frac{1}{N} \log p(x) &= \frac{1}{N} \log \left\{ \sum_{|\alpha|=N} a_\alpha x^\alpha \right\} \\ &\geq \frac{1}{N} \sum_{|\alpha|=N} a_\alpha \log(x^\alpha) \quad \text{by concavity of } \log, \text{ and (1) and (2)} \\ &= \frac{1}{N} \sum_{|\alpha|=N} a_\alpha \{ \alpha_1 \log x_1 + \cdots + \alpha_n \log x_n \} \\ &= \frac{1}{N} \sum_{j=1}^n \left\{ \sum_{|\alpha|=N} a_\alpha \alpha_j \log x_j \right\} = \frac{k}{N} \sum_{j=1}^n \log x_j \quad \text{by (3)}, \\ &= \frac{k}{N} \log(x_1 \cdots x_n). \end{aligned}$$

The value of k is determined by Euler's Formula

$$\langle D_x p, x \rangle = Np(x),$$

followed by assumptions (3) and then (2), yielding

$$\langle D_e p, e \rangle = \langle ke, e \rangle = kn = Np(e) = N,$$

i.e.,

$$k = \frac{N}{n}.$$

Substituting $1/n$ for k/N in the inequality above, we have

$$\frac{1}{N} \log p(x) \geq \frac{1}{n} \log(x_1 \cdots x_n).$$

□

Corollary 2.2 Let $p(\lambda) = p(\lambda_1, \dots, \lambda_n) \not\equiv 0$ be a symmetric homogeneous (real) polynomial of degree N all of whose coefficients are ≥ 0 . Normalize p so that $p(1, \dots, 1) = 1$. Then

$$p(\lambda_1, \dots, \lambda_n)^{\frac{1}{N}} \geq (\lambda_1 \cdots \lambda_n)^{\frac{1}{n}} \tag{2.3}$$

for all $\lambda \in \mathbb{R}_{>0}^n$.

Proof We need to prove the Central Ray Hypothesis, that all the $\frac{\partial p}{\partial \lambda_j}(e)$ agree (and equal k). This follows from the invariance under permutations. Then, as noted above, $k = N/n > 0$.

□

This corollary can be restated on the space $\text{Sym}^2(\mathbb{R}^n)$ of second derivatives.

Theorem 2.3 Let F be a homogeneous polynomial of degree N on $\text{Sym}^2(\mathbb{R}^n)$ given by

$$F(A) \equiv p(\lambda_1, \dots, \lambda_n) \quad (2.4)$$

where p is as in the Corollary 2.2 and $\lambda_1, \dots, \lambda_n$ are the eigenvalues of A . Note that F is normalized so that $F(I) = 1$. Then for all $A \geq 0$.

$$F(A)^{\frac{1}{N}} \geq (\det A)^{\frac{1}{n}}. \quad (2.5)$$

Note 2.4 If we drop the normalization Assumption (2) on p , and then replace $p(x)$ by $p(x)/p(e)$ which satisfies (2), we have for all $A \geq 0$ that

$$F(A)^{\frac{1}{N}} \geq \gamma (\det A)^{\frac{1}{n}}, \quad \text{where } \gamma \equiv F(I)^{\frac{1}{N}} = p(e)^{\frac{1}{N}}. \quad (2.6)$$

We will apply Theorem 2.3 to real symmetric, complex Hermitian symmetric, quaternionic Hermitian symmetric matrices A , and also the 2×2 octonian Hermitian symmetric matrices.

3 Gårding-Dirichlet polynomial operators

Let $F : \text{Sym}^2(\mathbb{R}^n) \rightarrow \mathbb{R}$ be a real homogeneous polynomial of degree N which is *Gårding hyperbolic with respect to the Identity I* . This mean that for each $A \in \text{Sym}^2(\mathbb{R}^n)$, the polynomial $t \mapsto F(tI + A)$ has all real roots. The *Gårding cone Γ* of F is the connected component of $\text{Sym}^2(\mathbb{R}^n) - \{F = 0\}$ which contains I . This is an open convex cone with 0 as the vertex. and for all directions $B \in \Gamma$, F is *B-hyperbolic*, which means that the polynomial $t \mapsto F(tB + A)$ has all real roots for every $A \in \text{Sym}^2(\mathbb{R}^n)$. We always assume that $F(I) > 0$. For complete details, see [8, 16, 17].

Definition 3.1 We assume a *positivity hypothesis*: $\Gamma \supset \{A : A > 0\}$. This defines a *Gårding-Dirichlet polynomial*, and applying this to the second derivative D^2u of functions u in \mathbb{R}^n gives a *Gårding-Dirichlet (or G-D) operator*. We shall always discuss a G-D operator in terms of the G-D polynomial that defines it. So the terms “operator” and “polynomial” will be used interchangeably here.

These operators have a form of strict ellipticity on the Gårding cone Γ :

$$F(A + P) > F(A) \quad \text{for all } A \in \Gamma \text{ and } P > 0. \quad (3.1)$$

Recall that $F^{\frac{1}{N}}$ is concave in Γ (see [8] or (D.2) below), which implies

$$F(A + P)^{\frac{1}{N}} \geq F(A)^{\frac{1}{N}} + F(P)^{\frac{1}{N}} \quad \text{for all } A \in \Gamma \text{ and } P > 0. \quad (3.2)$$

Combining this with (2.5) gives a stronger form of (3.1):

$$F(A + P)^{\frac{1}{N}} - F(A)^{\frac{1}{N}} \geq (\det P)^{\frac{1}{n}} \quad \text{for all } A \in \Gamma \text{ and } P > 0. \quad (3.3)$$

where $F(I) = 1$. Letting $A \rightarrow 0$ in (3.3) yields (2.5), so (3.3) and (2.5) are equivalent [2].

Basic G-D operators include the elementary symmetric functions $\sigma_k(A)$ (or Hessian operators). Two new ones are the p -fold sum operator (3.5) below, and the Lagrangian Monge-Ampère operator mentioned above (see [16] for these and many others including for example the δ -uniformly elliptic operators $Q_\delta(A) = \prod_j (\lambda_j(A) + \delta \text{tr} \lambda(A))$).

We also have the following construction.

Proposition 3.2 *Let F be a G-D operator of degree N . Then for any $P > 0$ the operator*

$$\delta_P F(A) \equiv \frac{d}{dt} F(tP + A) \Big|_{t=0} = \langle (\nabla F)_A, P \rangle$$

is a G-D operator of degree $N - 1$.

Proof The zeros of $\varphi'(t)$ interlace the N real zeros of $\varphi(t) \equiv F(tP + A)$, so $\delta_P F$ is Gårding. This also implies that the Gårding cones satisfy $\Gamma_{\delta_P F} \supset \Gamma_F$ (see [8]), and $\Gamma_F \supset \{A : A > 0\}$, so $\delta_P F$ satisfies positivity. \square

This construction can be iterated. For a G-D operator F , let $F(tI + A) = \prod_{j=1}^N (t + \lambda_j^F(A))$ where $\lambda_j^F(A)$ are called the *eigenvalues* of F with respect to I . Then differentiating this product at $t = 0$ and applying Proposition 3.2 yields that

$$\frac{1}{k!} \delta_I^k F(A) = \sigma_{N-k}(\lambda_1^F(A), \dots, \lambda_N^F(A)) \quad (3.4)$$

for $k = 0, 1, \dots, N$ are G-D polynomials, where σ_j is the j^{th} elementary symmetric function. It follows that *the elementary symmetric functions of $\lambda_1^F(A), \dots, \lambda_N^F(A)$ are polynomials in A .*

The next example was introduced in [14], along with p -geometry/ p -potential theory.

Example 3.3 (The p -Fold Sum Operator) Let

$$F(A) = \prod_{|J|=p} \lambda_J(A) \quad (3.5)$$

where

$$\lambda_J(A) = \lambda_{j_1}(A) + \dots + \lambda_{j_p}(A) \quad \text{where } j_1 < \dots < j_p \text{ and degree } = N = \binom{n}{p}.$$

(The inequality (2.5) for this example was first proven in [7].)

We shall see in §7 that the family of invariant G-D polynomials is huge! Concerning ellipticity.

Proposition 3.4 *A complete G-D operator F has uniformly elliptic linearization at each point of its Gårding cone Γ . That is, at $A \in \Gamma$*

$$B \mapsto \langle (\nabla F)_A, B \rangle \text{ has coefficient matrix } (\nabla F)_A > 0.$$

Proof This is Proposition B.1 in Appendix B. \square

Note that the uniform ellipticity $(\nabla F)_A > 0$ is stronger than $\langle (\nabla F)_A, P \rangle > 0, \forall P > 0$ mentioned earlier, and it requires the completeness hypothesis. Completeness means that all the variables in \mathbb{R}^n are needed to define F . This rules out operators such as $F(D^2u) = \partial^2 u / \partial x_1^2$ ($n \geq 2$). The invariant operators in the next section are always complete.

4 Invariant G-D operators

We shall be concerned principally with three cases.

Case 1 (Real Invariant G-D Operators) A G-D polynomial F on $\text{Sym}^2(\mathbb{R}^n)$ of degree N is real invariant if

$$F(A) = F(gAg^t) \quad \text{for all } g \in \text{O}(n).$$

Every such polynomial canonically determines an operator on every riemannian manifold.

By Theorem 4.1 below, F is *real invariant* if and only if

$$F(A) = p(\lambda_1(A), \dots, \lambda_n(A))$$

where p is a homogeneous symmetric polynomial of degree N in the eigenvalues of A . This polynomial $p(\lambda_1, \dots, \lambda_n)$ on \mathbb{R}^n satisfies the Assumptions (1) and (3) of the Basic Lemma 2.1, and $p(e) = F(I) > 0$.

Case 2 (Complex Invariant G-D Operators) Let $\text{Sym}_{\mathbb{C}}^2(\mathbb{C}^n)$ be the space of Hermitian symmetric $n \times n$ -matrices on \mathbb{C}^n . A (real-valued) G-D polynomial F on $\text{Sym}_{\mathbb{C}}^2(\mathbb{C}^n)$ of degree N is *complex invariant* if

$$F(A_{\mathbb{C}}) = F(gA_{\mathbb{C}}g^t) \quad \text{for all } g \in \text{U}(n).$$

Every such polynomial F canonically determines an operator on every Hermitian complex (or almost complex) manifold.

By Theorem 4.1 below, F is complex invariant if and only if

$$F(A_{\mathbb{C}}) = p(\lambda_1(A_{\mathbb{C}}), \dots, \lambda_n(A_{\mathbb{C}}))$$

where p is a homogeneous symmetric polynomial of degree N in the eigenvalues of $A_{\mathbb{C}}$. This polynomial $p(\lambda_1, \dots, \lambda_n)$ on \mathbb{R}^n satisfies the Assumptions (1) and (3) of the Basic Lemma 2.1, and $p(e) = F(I) > 0$.

This case can be looked at from the real point of view. Let $\mathbb{C}^n = (\mathbb{R}^{2n}, J)$. Then $\text{Sym}_{\mathbb{C}}^2(\mathbb{C}^n)$ can be identified with the subset of $A \in \text{Sym}^2(\mathbb{R}^{2n})$ satisfying $AJ = JA$. There is a projection $(\cdot)_{\mathbb{C}} : \text{Sym}^2(\mathbb{R}^{2n}) \rightarrow \text{Sym}_{\mathbb{C}}^2(\mathbb{C}^n)$ given by setting

$$A_{\mathbb{C}} = \frac{1}{2}(A - JAJ)$$

The real eigenvalues of $A_{\mathbb{C}}$ are the eigenvalues of $A_{\mathbb{C}}$ viewed as a Hermitian symmetric matrix, but where each eigenvalue now appears twice.

Case 3 (Quaternionic Invariant G-D Operators) Here we take the real point of view. Let $\mathbb{H}^n = (\mathbb{R}^{4n}, I, J, K)$ where I, J, K are complex structures satisfying the usual relations. (Thus \mathbb{H}^n can be viewed as a right quaternion vector space with scalars $\alpha + \beta I + \gamma J + \delta K$.) Then $\text{Sym}_{\mathbb{H}}^2(\mathbb{H}^n)$, the space of quaternionic Hermitian symmetric $n \times n$ -matrices, can be considered as the subset of $A \in \text{Sym}^2(\mathbb{R}^{4n})$ which commute with I, J and K .

There is a projection $(\cdot)_{\mathbb{H}} : \text{Sym}^2(\mathbb{R}^{4n}) \rightarrow \text{Sym}_{\mathbb{H}}^2(\mathbb{H}^n)$ given by setting

$$A_{\mathbb{H}} = \frac{1}{4}(A - IAI - JAJ - KAK).$$

Using the natural inner product on $\text{Sym}^2(\mathbb{R}^{4n})$, this map from A to $A_{\mathbb{H}}$ is orthogonal projection. Considering $A_{\mathbb{H}} \in \text{Sym}_{\mathbb{H}}^2(\mathbb{H}^n)$ as an element of $\text{Sym}^2(\mathbb{R}^{4n})$, it has a canonical form under the conjugate action of $\text{O}(4n)$, namely

$$A_{\mathbb{H}} = \lambda_1 P_{W_1} + \dots + \lambda_s P_{W_s} \quad (4.1)$$

with $\lambda_1 < \lambda_2 < \dots < \lambda_s$ and $\mathbb{R}^{4n} = W_1 \oplus \dots \oplus W_s$ an orthogonal decomposition, comprising the distinct eigenvalues and corresponding eigenspaces. Here P_W denotes orthogonal projection onto the real subspace $W \subset \mathbb{R}^{4n}$. Moreover, if v is an eigenvector of $A_{\mathbb{H}} \in \text{Sym}^2(\mathbb{R}^{4n})$ with eigenvalue λ , then vI, vJ, vK are also eigenvectors with the same eigenvalue λ . (Here vQ means right multiplication by the quaternion q .) That is,

$$\text{each eigenspace } W_j \text{ is a right quaternion vector subspace of } \mathbb{H}^n. \quad (4.2)$$

Taken together, (4.1) and (4.2) provide the *canonical form* for $A_{\mathbb{H}} \in \text{Sym}_{\mathbb{H}}^2(\mathbb{H}^n)$ under the conjugate action of $\text{Sp}(n) \cdot \text{Sp}(1) \equiv \text{Sp}(n) \times_{\mathbb{Z}_2} \text{Sp}(1)$. (Note that this canonical form could also be expressed using the eigenvalues $\lambda_j^0(A_{\mathbb{H}})$, $j = 1, \dots, n$, listed to multiplicity.)

A homogeneous real-valued polynomial F on $\text{Sym}_{\mathbb{H}}^2(\mathbb{H}^n)$ is G-D if $t \mapsto F(tI + A_{\mathbb{H}})$ has all real roots for every $A_{\mathbb{H}} \in \text{Sym}_{\mathbb{H}}^2(\mathbb{H}^n)$.

Let F be a real homogeneous polynomial F on $\text{Sym}_{\mathbb{H}}^2(\mathbb{H}^n)$ of degree N . Then F is *quaternionic invariant* if

$$F(A_{\mathbb{H}}) = F(gA_{\mathbb{H}}g^t) \quad \text{for all } g \in \text{Sp}(n) \cdot \text{Sp}(1).$$

Every such polynomial canonically determines an operator on every manifold with a topological \mathbb{H} -structure and a compatible metric, in particular a hyperKähler manifold with a Calabi-Yau metric.

By Theorem 4.1 below, a G-D polynomial F of degree N on $\text{Sym}_{\mathbb{H}}^2(\mathbb{H}^n)$ is quaternionic invariant if and only if

$$F(A_{\mathbb{H}}) = p(\lambda_1^0(A_{\mathbb{H}}), \dots, \lambda_n^0(A_{\mathbb{H}})) \quad (4.3)$$

where p is a homogeneous symmetric polynomial of degree N , and $\lambda_j^0(A_{\mathbb{H}})$ are the eigenvalues of $A_{\mathbb{H}}$ listed to multiplicity. This polynomial $p(\lambda_1, \dots, \lambda_n)$ on \mathbb{R}^n satisfies the Assumptions (1) and (3) of the Basic Lemma 2.1, and $p(e) = F(I) > 0$.

Theorem 4.1 *Let $F(A)$ be a real invariant G-D polynomial of degree N on $\text{Sym}^2(\mathbb{R}^n)$. Then*

$$F(A) = p(\lambda_1(A), \dots, \lambda_n(A))$$

where $p(\lambda_1, \dots, \lambda_n)$ is a symmetric homogeneous polynomial of degree N on \mathbb{R}^n which satisfies Assumptions (1) and (3) of Lemma 2.1 and for which $p(e) > 0$. Furthermore, p is G-D, that is p is Gårding hyperbolic w.r.t. e and the set $\{\lambda \in \mathbb{R}^n : \lambda_j > 0 \ \forall j\}$ is contained in its Gårding cone $\Gamma(p)$.

Conversely, if F is given this way for such a p , then F is a real invariant G-D operator.

The analogous statements hold in the complex and quaternionic cases.

Before the proof we make several remarks.

Remark 4.2 It is important to note in the three determinant cases:

$$(1) \ \det_{\mathbb{R}} A, \ A \in \text{Sym}_{\mathbb{R}}^2(\mathbb{R}^n), \quad (2) \ \det_{\mathbb{C}} A_{\mathbb{C}}, \ A \in \text{Sym}_{\mathbb{R}}^2(\mathbb{C}^n),$$

$$\text{or} \quad (3) \ \det_{\mathbb{H}} A_{\mathbb{H}}, \ A \in \text{Sym}_{\mathbb{R}}^2(\mathbb{H}^n),$$

that F is a G-D polynomial with Gårding I -eigenvalues given by

$$(1) \ \lambda(A) = \{\lambda_1(A), \dots, \lambda_n(A)\}, \quad (2) \ \lambda(A) = \{\lambda_1(A_{\mathbb{C}}), \dots, \lambda_n(A_{\mathbb{C}})\},$$

or (3) $\lambda(A) = \{\lambda_1(A_{\mathbb{H}}), \dots, \lambda_n(A_{\mathbb{H}})\}$, respectively,

defined in each case by the appropriate canonical form.

Remark 4.3 The fact that $F(A) \equiv \det_{\mathbb{H}} A_{\mathbb{H}} \equiv \lambda_1(A_{\mathbb{H}}) \cdots \lambda_n(A_{\mathbb{H}})$ is a polynomial in $A \in \text{Sym}_{\mathbb{R}}^2(\mathbb{H}^n)$ (or equivalently a polynomial in $A_{\mathbb{H}} \in \text{Sym}_{\mathbb{H}}^2(\mathbb{H}^n) \subset \text{Sym}_{\mathbb{R}}^2(\mathbb{R}^{4n})$) requires proof. The Moore determinant provides an algebraic construction which is rather complicated (see [3].) Some might prefer the following proof.

Lemma 4.4 *The determinant $\det_{\mathbb{H}} A_{\mathbb{H}} \equiv \lambda_1(A_{\mathbb{H}}) \cdots \lambda_n(A_{\mathbb{H}})$ is a polynomial in $A \in \text{Sym}_{\mathbb{R}}^2(\mathbb{H}^n)$.*

Proof Let $\mathbb{H} = (\mathbb{R}^{4n}, I, J, K)$ and let $\pi : \text{Sym}^2(\mathbb{R}^{4n}) \rightarrow \text{Sym}^2(\mathbb{R}^{4n})$ be given by

$$\pi(A) = \frac{1}{4}(A - IAI - JAJ - KAK) \equiv A_{\mathbb{H}}.$$

Consider

$$\begin{aligned} \det(I + tA_{\mathbb{H}}) &= 1 + \Sigma_1(A)t + \Sigma_2(A)t^2 + \cdots + \Sigma_{4n}t^{4n} \\ &= (1 + \sigma_1(\lambda)t + \sigma_2(\lambda)t^2 + \sigma_3(\lambda)t^3 + \cdots + \sigma_n(\lambda)t^n)^4 \end{aligned}$$

where the λ 's are the n eigenvalues of $A_{\mathbb{H}}$. Now $\det(I + tA_{\mathbb{H}})$ is a polynomial in (A, t) so the $\Sigma_k(A)$'s are polynomials in A . We also have

$$\begin{aligned} &(1 + \sigma_1(\lambda)t + \sigma_2(\lambda)t^2 + \sigma_3(\lambda)t^3 + \cdots)(1 + \sigma_1(\lambda)t + \sigma_2(\lambda)t^2 + \sigma_3(\lambda)t^3 + \cdots) \\ &\cdot (1 + \sigma_1(\lambda)t + \sigma_2(\lambda)t^2 + \sigma_3(\lambda)t^3 + \cdots)(1 + \sigma_1(\lambda)t + \sigma_2(\lambda)t^2 + \sigma_3(\lambda)t^3 + \cdots) \\ &= 1 + 4\sigma_1(\lambda)t + [4\sigma_2(\lambda) + 6\sigma_1^2(\lambda)]t^2 + [4\sigma_3(\lambda) + \cdots]t^3 + [4\sigma_4(\lambda) + \cdots]t^4 + \cdots \\ &= 1 + \Sigma_1(A)t + \Sigma_2(A)t^2 + \Sigma_3(A)t^3 + \Sigma_4(A)t^4 + \cdots \end{aligned}$$

Now

$4\sigma_1(\lambda) = \Sigma_1(A)$ is a polynomial in A . Therefore

$4\sigma_2(\lambda) + 6\sigma_1^2(\lambda) = \Sigma_2(A)$ is a polynomial in A . Hence $\sigma_2(\lambda)$ is a polynomial in A .

In general $4\sigma_k(\lambda) + \text{a polynomial in } A = \Sigma_k(A)$ and so $\sigma_k(\lambda)$ is a polynomial in A . \square

A polynomial $p(\lambda)$ which satisfies all the conditions in Theorem 4.1 is defined (Def. 7.3) to be a universal G-D polynomial.

Theorem 4.1 is generalized in Theorem 7.2. There the Gårding I -eigenvalues used in Theorem 4.1 for one of the three determinants $\det_{\mathbb{R}}$, $\det_{\mathbb{C}}$, $\det_{\mathbb{H}}$ are replaced by the Gårding I -eigenvalues of any G-D operator.

Proof of Theorem 4.1 We begin by restricting F to the subspace $\mathbb{D} \subset \text{Sym}^2(\mathbb{R}^n)$ of diagonal matrices. For each $\lambda \in \mathbb{R}^n$, let $M(\lambda) = \sum_i \lambda_i e_i \circ e_i$ denote the diagonal matrix with entries $\lambda_1, \dots, \lambda_n$. Define

$$p(\lambda) \equiv F(M(\lambda)) \quad \text{for } \lambda \in \mathbb{R}^n.$$

Then $p(\lambda)$ is a homogenous polynomial of degree N . The fact that F is invariant under the orthogonal group implies that p is invariant under permutations of the λ_j 's, i.e., p is a symmetric polynomial. Now by the conjugation invariance, this polynomial determines F by the formula $F(A) = p(\lambda_1(A), \dots, \lambda_n(A))$. So now everything is reconstructed in terms of p . The positivity hypothesis that $F > 0$ on $\{A : A > 0\}$ is equivalent to the strict positivity of p on $\mathbb{R}_{>0}^n$, since $A > 0 \iff \lambda(A) \in \mathbb{R}_{>0}^n$. Assertion (1) that all the coefficients of p are ≥ 0 , can be proved as follows.

Note that

$$\frac{1}{\alpha_1! \cdots \alpha_n!} \frac{\partial^{|\alpha|} p}{\partial x_1^{\alpha_1} \cdots \partial x_n^{\alpha_n}}$$

equals the coefficient a_α for each multi-index α of length $|\alpha| \equiv \sum_k \alpha_k = N$. To prove that this α^{th} partial of p is ≥ 0 we use Proposition 3.2.

Since $p(\lambda) = F(\lambda_1 e_1 \circ e_1 + \cdots + \lambda_n e_n \circ e_n)$, the partials of p

$$\frac{\partial p}{\partial \lambda_j}(\lambda) = \langle (\nabla F)_{M(\lambda)}, e_j \circ e_j \rangle$$

are equal to the directional derivatives of F in the directions $P_j = e_j \circ e_j \geq 0$. By Proposition 3.2 we have that $\langle (\nabla F)_A, P \rangle = (\delta_P F)(A) > 0$ for all $A \in \Gamma$ and $P > 0$. Hence, $(\delta_P F)(A) \geq 0$ for all $A \in \Gamma$ and $P \geq 0$.

By Proposition 3.2, $\delta_P F$ is also an invariant G-D operator. Therefore, this process can be repeated N times proving that the N^{th} directional derivatives in directions $P_1 > 0, \dots, P_N > 0$ are > 0 . Hence, the N^{th} directional derivatives in directions $P_1 \geq 0, \dots, P_N \geq 0$ are ≥ 0 . Taking these latter P_j 's to each be in axis directions $e_k \circ e_k$ yields the desired result that the α^{th} partial of p , $|\alpha| = N$, is ≥ 0 . So Assertion (1) is proved.

For Assumption (3) we use the fact that the gradient $D_e p$ at e is invariant under permutation of the λ_j 's since $e = (1, \dots, 1)$ is invariant and so is p . Since $\mathbb{R}e$ is the only subspace of \mathbb{R}^n fixed under the permutation group, we have $D_e p = k e$ for some $k \in \mathbb{R}$. However k must be positive since by Euler's formula $nk = \langle D_e p, e \rangle = Np(e) > 0$ as in the proof of Basic Lemma 2.1.

The arguments for the complex and quaternionic cases are exactly the same. \square

Theorem 4.5 *Let F be an invariant G-D operator from one of the three cases above. Let $\det(A)$ denote the complex or quaternionic determinant in the complex and quaternionic cases. Then*

$$F(A)^{\frac{1}{N}} \geq F(I)^{\frac{1}{N}} \det(A)^{\frac{1}{n}} \quad (4.4)$$

Proof This follows directly from Theorem 4.1 and Basic Lemma 2.1. \square

5 Relevance to the work of Guo-Phong-Tong and Guo-Phong

One of the principle motivations for this work was the recent paper of B. Guo, D. H. Phong, and F. Tong [10], which among other things gave a purely PDE proof of the C^0 -estimate in Yau's proof of the Calabi Conjecture. Their main theorem, Theorem 1, has much broader applicability to operators of complex invariant type (as in Sect. 3 in this paper) on Kähler manifolds. The main assumption (1.4) in this theorem follows if one can prove the majorization of the determinant formula in the cone $\text{Int } \mathcal{P}$ of positive definite matrices (see [[10], Lemma 4]). That is exactly what we have done for all complex invariant Gårding-Dirichlet operators, and, therefore, for all the associated operators on Kähler manifolds.

In a later paper of Guo and Phong [10], a number of important results for operators on Hermitian manifolds were established. Again the important assumption is established by the majorization of the determinant formula. Hence, all the results in this paper hold for every operator on a Hermitian manifold, which is induced by any complex invariant Gårding-Dirichlet operator.

6 Interior regularity-work of Abja-Olive

In this section we establish C^∞ interior regularity for $W^{2,p}$ -solutions of complex invariant G-D operators (in the Euclidean case). This is based on results of Abja-Olive [1]. We begin by showing that the basic assumptions (a)–(f) in [1] are satisfied. As above we work in eigenvalue space with a symmetric polynomial $p(\lambda_1, \dots, \lambda_n)$ which satisfies the assumptions of Lemma 2.1. Let Γ denote the connected component of $\mathbb{R}^n - \{p = 0\}$ containing e .

- (a) $\mathbb{R}_{>0}^n \subset \Gamma \subset \{\lambda_1 + \dots + \lambda_n > 0\}$.
- (b) p is a C^1 on $\bar{\Gamma}$.
- (c) $p > 0$ on Γ .
- (d) p is positively homogeneous of degree N on Γ .
- (e) $p^{\frac{1}{N}}$ is concave on Γ .
- (f) $p(\lambda)^{\frac{1}{N}} \geq (\lambda_1 \cdots \lambda_n)^{\frac{1}{n}}$ with equality on $\text{span}(e)$.

Proof The left hand inclusion of (a) follows from the positivity hypothesis. For the right hand inclusion, note that Γ is convex and invariant under permutations of the λ_j 's. Among half spaces H containing Γ with the vertex $0 \in \partial H$, there is only one H_0 which is invariant under permutations. Since $\mathbb{R} \cdot e$ is the only invariant line, we have $H_0 = e^\perp = \{\lambda_1 + \dots + \lambda_n \geq 0\}$.

Hypothesis (b) is clear since p is a polynomial. Hypothesis (c) follows from the definition of Γ . Hypothesis (d) is an assumption. Hypothesis (e) is Theorem 2 in [8] for the initial polynomial on matrices. This restricts to concavity for diagonal matrices. Hypothesis (f) is Corollary 2.2.

Now the assumptions of [1] are on matrix space and not on eigenvalue space \mathbb{R}^n . One needs to know that $C \subset \mathbb{R}^n$ is convex if and only if $\lambda^{-1}(C)$ is convex (where $\lambda^{-1}(C) = \text{all matrices whose vector of eigenvalues lies in } C$). This was proved in [4]. (See also [[17], Thm. 8.4] and [AO, Prop. 3.1].) One also needs to know that if $C \subset \mathbb{R}^n$ is compact, then so is $\lambda^{-1}(C)$. However, $C \times O(n)$ is compact and $\lambda^{-1}(C)$ is an image of $C \times O(n)$. \square

We can now apply the Main Theorem 1.1 in [1] where everything takes place in \mathbb{C}^n and the authors deal with Hermitian symmetric matrices. Let F be the invariant G-D operator on \mathbb{C}^n given by a homogeneous polynomial $p(\lambda_1, \dots, \lambda_n)$ of degree N as above, applied to the eigenvalues of Hermitian symmetric matrices A , that is, $F(A) = p(\lambda(A))$. We now let Γ denote the Gårding cone of F , and we fix a domain $\Omega \subset \mathbb{C}^n$.

Theorem 6.1 *Let $u \in W_{\text{loc}}^{2,q}(\Omega)$, $q > n \max\{N - 1, 1\}$, with $D^2u(x) \in \Gamma$ for a.e. $x \in \Omega$ satisfying*

$$F(D^2u) = f \quad \text{a.e. in } \Omega$$

where $f > 0$ and in $C^2(\Omega)$. Then for each domain $\Omega_0 \subset\subset \Omega$ there exists $R > 0$ with

$$\|\Delta u\|_{L^\infty(\Omega_0)} \leq R.$$

We now apply this bound on Δu . We begin with the following.

(g) (Sets of Uniform Ellipticity) For each $R > 0$, F is uniformly elliptic in the region

$$\Gamma_R = \left\{ A \in \Gamma : \text{tr}(A) < R \text{ and } \frac{1}{R} < F(A) \right\}$$

Now the polynomial F has the property that for $A \in \Gamma$,

$$D_A F \in \Gamma^* \text{ (the “open polar” of } \Gamma) \subset \text{Int } \mathcal{P}.$$

The definition of the *open polar* Γ^* is in Appendix A. One sees easily that an invariant D-G polynomial satisfies the completeness condition given in Appendix A. Therefore, Proposition B.1 in Appendix B applies to show that $D_A F \in \Gamma^*$. Since $\text{Int } \mathcal{P} \subset \Gamma$, we have $\Gamma^* \subset \text{Int } \mathcal{P}$. Therefore the linearization of F is positive definite on all of Γ . Assumption (g) is then a result of the following fact.

(g)' For all $R > 0$ the set

$$\bar{\Gamma}_R \equiv \left\{ A \in \Gamma : \sum_{j=1}^n \lambda_j(A) \leq R \text{ and } \frac{1}{R} \leq F(A) \right\} \text{ is compact.}$$

Proof From (a) above one easily sees that $I \in \Gamma^0$. The assertion (g)' then follows from Corollary C.3 in Appendix C with $c_1 = R$, $c_2 = \log(1/R)$ and $\mathfrak{g} = F$. \square

The fact that (g) holds in this generality is quite useful, so we state it separately.

Proposition 6.2 *For each $R > 0$, the G-D operator F restricted to the set $\bar{\Gamma}_R$ is uniformly elliptic.*

In particular, we have interior regularity for $W^{2,p}$ solutions (see [1, 5, 20]).

Theorem 6.3 *Let F be an invariant complex G-D operator. Under the assumptions of Theorem 6.1,*

$$f \in C^\infty(\Omega) \Rightarrow u \in C^\infty(\Omega).$$

7 On the enormous universe of invariant G-D operators

Let's fix one of the algebras \mathbb{R} , \mathbb{C} or \mathbb{H} , and for the discussion in this chapter “invariant” will always mean “real invariant”, “complex invariant” or “quaternionic invariant” accordingly. The discussion is identical in these three cases.

We begin with the following.

Proposition 7.1 *The product of two invariant G-D polynomials is again an invariant G-D polynomial, with eigenvalue set the union of the F and G eigenvalues.*

Proof If

$$F(tI + A) = F(I) \prod_{k=1}^n (t + \lambda_k^F(A)) \text{ and } G(tI + A) = G(I) \prod_{k=1}^n (t + \lambda_k^G(A)),$$

then

$$F(tI + A)G(tI + A) = F(I)G(I) \prod_{k=1}^n (t + \lambda_k^F(A)) \prod_{k=1}^n (t + \lambda_k^G(A)),$$

and so FG is Gårding hyperbolic w.r.t. I . If F and G are each invariant under a group \mathcal{G} acting on the A 's, so is the product. \square

Proposition 7.2 *Let F be an invariant G-D polynomial of degree N . Then the operator*

$$(\delta_I F)(A) \equiv \left. \frac{d}{dt} F(tI + A) \right|_{t=0} = \langle (\nabla F)_A, I \rangle \quad (7.1)$$

is an invariant G-D polynomial of degree $N - 1$.

Proof The invariance of $\delta_I F$ is clear. The rest is Proposition 3.2. \square

We formalize the definition given after Theorem 4.1, which has been taken from [16].

Definition 7.3 A *universal G-D polynomial* is a homogeneous symmetric polynomial $p(\lambda_1, \dots, \lambda_n)$ which is Gårding hyperbolic w.r.t. e and satisfies Assumptions (1) and (3) of Basic Lemma 2.1 with $p(e) > 0$. Note that Assumption (1) implies $p > 0$ on $\mathbb{R}_{>0}^n$.

These are exactly the polynomials such that $F(A) = p(\lambda^F(A))$ for invariant G-D polynomials F (Theorem 4.1).

Consider now a G-D polynomial F of degree M on $\text{Sym}^2(\mathbb{R}^n)$ with I-eigenvalues $\lambda^F(A) = (\lambda_1^F(A), \dots, \lambda_M^F(A))$. We shall construct a new G-D polynomial by applying a universal Gårding polynomial p of degree N in M variables to the F -eigenvalues. This is why p is called universal.

Theorem 7.4 Let p and F be as above. Define

$$P_F(A) \equiv p(\lambda_1^F(A), \dots, \lambda_M^F(A)) = p(\lambda^F(A))$$

for $A \in \text{Sym}^2(\mathbb{R}^n)$. Then P_F is also a G-D polynomial of degree N on $\text{Sym}^2(\mathbb{R}^n)$, with Gårding eigenvalues

$$\lambda_j^{P_F}(A) = \Lambda_j(\lambda^F(A)) \quad j = 1, \dots, N, \quad (7.2)$$

where $\Lambda_1(\lambda), \dots, \Lambda_N(\lambda)$, are the e -eigenvalues of the G-D polynomial p .

The Gårding cones are related by $\Gamma_{P_F} = (\lambda^F)^{-1}(\Gamma_p)$.

Furthermore, if F is invariant, so is P_F .

Proof We first observe that $P_F(A)$ is a polynomial. This is evident if $p(\lambda) = \lambda_1 \cdots \lambda_M$ where $P_F = F$. It then follows for all the elementary symmetric functions $\sigma_1, \dots, \sigma_M$ by Proposition 3.2 and Example 3.3. Now by a classical result, $p(\lambda)$ is a polynomial in the elementary symmetric functions, and so $P_F(A)$ is a polynomial.

Note first that $P_F(I) = p(\lambda^F(I)) = p(e) > 0$. By (2.1) in [8],

$$p(te + \lambda^F) = p(e) \prod_{j=1}^N (t + \Lambda_j(\lambda^F)).$$

Now $\lambda^F(I) = e$ since $F(tI + I) = (t + 1)^M F(I)$. Therefore (2)' in [8] says that

$$\lambda^F(tI + A) = te + \lambda^F(A),$$

and so

$$\begin{aligned} P_F(tI + A) &= p(\lambda^F(tI + A)) \\ &= p(te + \lambda^F(A)) \\ &= p(e) \prod_{j=1}^N (t + \Lambda_j(\lambda^F(A))). \end{aligned}$$

Therefore, $\Lambda_j(\lambda^F(A))$, $j = 1, \dots, N$, are the Gårding I -eigenvalues of P_F . The statement about Gårding cones follows from (7.2). The invariance statement follows from Theorem 4.1. \square

Many basic invariant G-D polynomials are given in [16, 17]. Taking these and using Propositions 7.2, 7.3, and 7.4 successively in long chains show that *the set of invariant G-D polynomials operators is very large*.

8 Determinant majorization for the Lagrangian Monge-Ampère operator

The Lagrangian Monge-Ampère Operator is a $U(n)$ -invariant G-D operator on \mathbb{C}^n which is dependent on the Skew part and independent of the traceless Hermitian symmetric part of a matrix A . Its pointwise linearizations are elliptic. For a compete discussion, including its importance for a lagrangian potential theory and its geometrical interpretation, one should see [18], where the operator was introduced. Note that by its $U(n)$ -invariance this operator is defined on any (almost) complex Hermitian manifold.

For the algebra pertaining to this equation, we shall be working on $\mathbb{C}^n = (\mathbb{R}^{2n}, J)$. The operator is defined on the $U(n)$ -invariant subspace $\mathbb{R}I \oplus \text{Herm}^{\text{Skew}}(\mathbb{C}^n) \subset \text{Sym}_{\mathbb{R}}^2(\mathbb{R}^{2n})$ where $\text{Herm}^{\text{Skew}}(\mathbb{C}^n) = \{A : JA = -AJ\}$. It is zero on the space of traceless Hermitian symmetric matrices (those traceless A with $JA = AJ$). Therefore we can assume that this part of A is zero. If $Ae = \lambda e$ and A is skew Hermitian, then $A(Je) = -\lambda Je$. Hence, we can let $\pm\lambda_1, \dots, \pm\lambda_n$ denote the eigenvalues of the skew part $A^{\text{skew}} \equiv \frac{1}{2}(A + JAJ)$. Then

$$A = tI + A^{\text{skew}}$$

with eigenvalues

$$t \pm \lambda_1, \dots, t \pm \lambda_n \quad \text{and} \quad \text{tr}(A) = 2nt.$$

Thus we have

$$\det(A) = \prod_{j=1}^n (t + \lambda_j)(t - \lambda_j).$$

With $\mu \equiv \frac{1}{2}\text{tr}(A) = nt$ the operator is defined in [18] by

$$\begin{aligned} F(A) &= \prod_{2^n \pm} (\mu \pm \lambda_1 \pm \lambda_2 \pm \dots \pm \lambda_n) = \prod_{2^n \pm} (nt \pm \lambda_1 \pm \lambda_2 \pm \dots \pm \lambda_n) \\ &= \prod_{2^n \pm} (e_1^{\pm} + \dots + e_n^{\pm}) \quad \text{where } e_j^{\pm} = t \pm \lambda_j. \end{aligned}$$

The eigenvalues of A are e_j^+ and e_j^- for $j = 1, \dots, n$, so

$$\det(A) = e_1^+ e_1^- \cdots e_n^+ e_n^-$$

Proposition 8.1 *In the region where $A > 0$,*

$$F(A)^{\frac{1}{2^n}} \geq \det(A)^{\frac{1}{2^n}}$$

Proof Positivity implies that $e_j^{\pm} > 0$ for all j . Hence, for all 2^{n-1} choices of \pm we have

$$\log(e_1^{\pm} + \dots + e_j^+ + \dots + e_n^{\pm}) + \log(e_1^{\pm} + \dots + e_j^- + \dots + e_n^{\pm}) \geq \log(e_j^+) + \log(e_j^-).$$

Summing over all 2^{n-1} choices gives

$$\begin{aligned} &\frac{1}{2^{n-1}} \sum_{2^{n-1} \pm} \left\{ \log(e_1^{\pm} + \dots + e_j^+ + \dots + e_n^{\pm}) + \log(e_1^{\pm} + \dots + e_j^- + \dots + e_n^{\pm}) \right\} \\ &= \frac{1}{2^{n-1}} \sum_{2^n \pm} \left\{ \log(e_1^{\pm} + \dots + e_j^{\pm} + \dots + e_n^{\pm}) \right\} = \frac{1}{2^{n-1}} \log F(A) \\ &\geq \log(e_j^+) + \log(e_j^-). \end{aligned}$$

Summing over $j = 1, \dots, n$ gives

$$\frac{n}{2^{n-1}} \log F(A) \geq \sum_{j=1}^n \{\log(e_j^+) + \log(e_j^-)\} = \log(\det(A))$$

□

9 Being a Gårding-Dirichlet operator is not enough for determinant majorization

It is not true that the determinant majorization inequality holds for all G-D polynomials. The Central Ray Hypothesis is crucial. For a counterexample we take $F : \text{Sym}^2(\mathbb{R}^2) \rightarrow \mathbb{R}$ to be the cubic polynomial

$$F(A) \equiv a_{11}^2 a_{22}.$$

The polynomial F is I -hyperbolic with Gårding eigenvalues $\lambda_1^F(A) = \lambda_2^F(A) = a_{11}$ and $\lambda_3^F(A) = a_{22}$ since

$$F(tI + A) = (t + a_{11})^2(t + a_{22}).$$

The Gårding cone Γ equals $\{A : a_{11} > 0 \text{ and } a_{22} > 0\}$, and since its closure $\overline{\Gamma}$ contains \mathcal{P} , this proves that F is a G-D operator.

Suppose now that $A > 0$ is diagonal, so that, $a_{12} = 0$. Then

$$\frac{F(A)^{\frac{1}{3}}}{\det(A)^{\frac{1}{2}}} = \left(\frac{a_{11}}{a_{22}} \right)^{\frac{1}{6}} \not\geq \gamma \quad \text{for any } \gamma > 0.$$

Of course the Central Ray Hypothesis is not satisfied by the diagonal operator $F(a_{11}, 0, a_{22})$, so the Basic Lemma 2.1 does not apply.

We note that since F is a G-D operator, the inhomogeneous Dirichlet problem

$$F(D^2u) = \left(\frac{\partial^2 u}{\partial x^2} \right)^2 \left(\frac{\partial^2 u}{\partial y^2} \right) = f(x, y), \quad \frac{\partial^2 u}{\partial x^2}, \frac{\partial^2 u}{\partial y^2} \geq 0, \quad u|_{\partial\Omega} = \varphi,$$

with $f \in C(\overline{\Omega})$, $f \geq 0$, and $\varphi \in C(\partial\Omega)$, can be uniquely solved on domains $\Omega \subset\subset \mathbb{R}^2$ with smooth strictly convex boundaries (see [19]).

This simple example extends to a family where the determinant majorization fails.

Consider the diagonal G-D operator

$$F(A) \equiv a_{11}^{N-1} \cdot \frac{1}{n} (a_{22} + \dots + a_{n+1,n+1}),$$

with $N \geq 3$, $n \geq 1$, on $\mathbb{R} \times \mathbb{R}^n$ with coordinates (t, x) . The Gårding cone Γ equals $\{a_{11} > 0 \text{ and } a_{22} + \dots + a_{n+1,n+1} > 0\}$. Restricting to diagonal $A > 0$ and fixing $a_{22}, \dots, a_{n+1,n+1}$, one has

$$\frac{F(A)^{\frac{1}{N}}}{(\det A)^{\frac{1}{n+1}}} = a_{11}^{1-\frac{1}{N}-\frac{1}{n+1}} C \quad \text{with } C > 0.$$

Since $1 - \frac{1}{N} - \frac{1}{n+1} \geq 1 - \frac{1}{3} - \frac{1}{2} = \frac{1}{6} > 0$, there is no lower bound for the left-hand term on $A > 0$, i.e., determinant majorization fails.

Although, again by [19], the continuous inhomogeneous Dirichlet Problem can be solved as above.

When $n \geq 2$, there exists a solution to $F(D^2u) = k > 0$ which is $\frac{2}{N}$ -Hölder continuous but no better. Note that the domain of the operator F is the space of u.s.c. functions $u(t, x)$ which are convex in t and Δ subharmonic in x .

Lemma 9.1 ($n \geq 2$). *The singular Pogorelov function $u(t, x) \equiv g(t)|x|^{\frac{2}{N}}$, where g satisfies $g''(t)^{N-1}g(t) = 1$, $g(0) = 1$, $g'(0) = 0$, is a viscosity solution to*

$$F(D^2u) \equiv u_{tt}^{N-1} \frac{1}{n} \Delta_x u = k \equiv \frac{2}{nN} \left(n - 2 + \frac{2}{N} \right) > 0. \quad (9.1)$$

Outline of Proof First, if $x \neq 0$, then using $\frac{\partial}{\partial x_j} |x| = \frac{x_j}{|x|}$ one directly computes that $\frac{1}{n} \Delta_x |x|^{\frac{2}{N}} = \frac{2}{nN} (n - 2 + 2/N) |x|^{\frac{2}{N}-2}$. Hence,

$$F(D^2u) = g''(t)^{N-1} g(t) \frac{1}{n} \Delta_x |x|^{\frac{2}{N}} = k.$$

Second, use the smooth approximations

$$u_\epsilon(t, x) \equiv g(t)(|x|^2 + \epsilon)^{\frac{1}{N}} \quad (9.2)$$

which decrease to u as $\epsilon \searrow 0$.

A similar direct calculation yields

$$F(D^2u_\epsilon) = \left(\frac{2}{nN} \right) \frac{C|x|^2 + \epsilon n}{|x|^2 + \epsilon}, \quad \text{with } C \equiv n - 2 + \frac{2}{N}. \quad (9.3)$$

Note that $k = \frac{2}{nN}C$ and that

$$\frac{C|x|^2 + \epsilon n}{|x|^2 + \epsilon} - C = \frac{\epsilon(n - C)}{|x|^2 + \epsilon}, \quad \text{with } n - C \equiv 2 - \frac{2}{N} \geq 1.$$

Hence, each u_ϵ is a subsolution of $F(D^2u) = k$, proving that the decreasing limit u is also a subsolution.

Finally, with $\epsilon > 0$ sufficiently small, one can show that there exist $\eta_\epsilon > 0$ with $\eta_\epsilon \rightarrow 0$ and $F(D^2(u_\epsilon - \eta_\epsilon \frac{1}{2}|x|^2)) \leq k$ if $u_\epsilon - \eta_\epsilon \frac{1}{2}|x|^2$ is admissible at x , i.e., $\Delta_x u_\epsilon - \eta_\epsilon \geq 0$. That is, $u_\epsilon - \eta_\epsilon \frac{1}{2}|x|^2$ is a supersolution of the equation $F(D^2u) = k$. As $\epsilon \rightarrow 0$, this converges uniformly to u . So u is also supersolution and therefore a viscosity solution. \square

10 Other applications of the Basic Lemma

There are other families of polynomial operators on $\text{Sym}^2(\mathbb{R}^n)$, besides those discussed above, to which the Basic Lemma 2.1 does apply.

Throughout this section we assume that $p(x)$ is a polynomial satisfying the Hypotheses of Basic Lemma 2.1 and that $E \subset \text{Sym}^2(\mathbb{R}^n)$ is a closed $\mathbb{R}_{\geq 0}^n$ -monotone set containing $\mathbb{R}_{\geq 0}^n$. We also assume the pair p, E satisfies

$$p(x + y) \geq p(x) \quad \forall x \in E \text{ and } y \in \mathbb{R}_{\geq 0}^n. \quad (10.1)$$

Since p has coefficients ≥ 0 (Assumption (1)), p always satisfies (10.1) if we set $E \equiv \mathbb{R}_{\geq 0}^n$.

FAMILY 1 (Diagonal Operators) Let F be a homogeneous polynomial of degree N on $\text{Sym}^2(\mathbb{R}^n)$ such that

$$F(A) = p(a_{11}, a_{22}, \dots, a_{nn}) \quad \text{if } (a_{11}, \dots, a_{nn}) \in E.$$

where p satisfies the conditions of Basic Lemma 2.1. Then F satisfies

$$F(A)^{\frac{1}{N}} \geq \det(A)^{\frac{1}{n}} \quad \text{for all } A > 0. \quad (10.2)$$

Note that F with domain restricted to $\{A \in \text{Sym}^2(\mathbb{R}^n) : (a_{11}, \dots, a_{nn}) \in E\}$ is an elliptic operator by (10.1).

Proof By the Basic Lemma 2.1, $F(A)^{\frac{1}{N}} \geq (a_{11}a_{22} \cdots a_{nn})^{\frac{1}{n}}$ for $A > 0$, and for $A > 0$ we have that $a_{11}a_{22} \cdots a_{nn} \geq \det(A)$. This last (well known) inequality is proved as follows. Take the Cholesky decomposition $A = LL'$ where L is lower triangular. Then $\det A = (\det L)(\det L') = (\det L)^2 = \alpha_1^2 \cdots \alpha_n^2$ where $\alpha_1, \dots, \alpha_n$ are the diagonal entries of L . In terms of row vectors we write L as

$$L = \begin{pmatrix} w_1 \\ \vdots \\ w_n \end{pmatrix} \quad \text{where} \quad \begin{aligned} w_1 &= (\alpha_1, 0, 0, \dots, 0) \\ w_2 &= (*, \alpha_2, 0, \dots, 0) \\ &\vdots \\ w_n &= (*, *, \dots, *, \alpha_n). \end{aligned}$$

Hence,

$$A = \begin{pmatrix} |w_1|^2 & & & \\ & |w_2|^2 & * & \\ & * & |w_n|^2 & \end{pmatrix}$$

and the diagonal entries are $a_{11} \cdots a_{nn} = |w_1|^2 |w_2|^2 \cdots |w_n|^2 \geq \alpha_1^2 \cdots \alpha_n^2$. \square

FAMILY 2 (Ordered Eigenvalue Operators) Let $p(\lambda_1, \dots, \lambda_n)$ satisfy the assumptions of Basic Lemma 2.1, and for any $A \in \text{Sym}^2(\mathbb{R}^n)$ set

$$F(A) \equiv p(\lambda_1(A), \dots, \lambda_n(A)) \quad \text{where } \lambda_1(A) \leq \lambda_2(A) \leq \cdots \leq \lambda_n(A).$$

Then F satisfies (10.2). (Note that these operators F are continuous but generally not polynomials.)

There are many non-symmetric polynomials satisfying these assumptions.

Lemma 10.1 Suppose $q(y_1, \dots, y_n)$ and $r(z_1, \dots, z_m)$ both satisfy the hypotheses of Basic Lemma 2.1, with the same value of k in Assumption (3), then so also does the polynomial

$$p(x_1, \dots, x_{n+m}) \equiv q(x_1, \dots, x_n) r(x_{n+1}, \dots, x_{n+m}).$$

Proof Note that $p(1, 1, \dots, 1) = q(1, \dots, 1) r(1, \dots, 1) = 1$. Also we have

$$\frac{\partial p}{\partial x_j}(1, \dots, 1) = \begin{cases} \frac{\partial q}{\partial x_j}(1, \dots, 1) r(1, \dots, 1) = k & \text{if } 1 \leq j \leq n \\ q(1, \dots, 1) \frac{\partial r}{\partial x_j}(1, \dots, 1) = k & \text{if } n+1 \leq j \leq n+m. \end{cases}$$

It is clear that the coefficients of p are all ≥ 0 .

Lemma 10.2 Suppose $q(y_1, \dots, y_n)$ and $r(z_1, \dots, z_m)$ both satisfy the hypotheses of Basic Lemma 2.1 and have the same degree. Let k and k' be the constants in Assumption (3) for q and r respectively. Then the polynomial

$$p(\lambda_1, \dots, \lambda_{n+m}) \equiv \frac{k'}{k+k'}q(\lambda_1, \dots, \lambda_n) + \frac{k}{k+k'}r(\lambda_{n+1}, \dots, \lambda_{n+m}).$$

also satisfies the hypotheses of Basic Lemma 2.1.

Proof We clearly have that p is homogeneous, and $p(1, \dots, 1) = 1$. If $j \leq n$,

$$\frac{\partial p}{\partial x_j}(1, \dots, 1) = \frac{k'}{k+k'} \frac{\partial q}{\partial x_j}(1, \dots, 1) = \frac{k'k}{k+k'},$$

and if $j \geq n+1$, then

$$\frac{\partial p}{\partial x_j}(1, \dots, 1) = \frac{k}{k+k'} \frac{\partial r}{\partial x_j}(1, \dots, 1) = \frac{k'k}{k+k'}.$$

Again it is clear that the coefficients of p are all ≥ 0 . \square

Example 10.3 The polynomials $P_j(x) = \frac{1}{j}\{x_1^j + \dots + x_n^j\}$ for $j \geq 1$ each satisfy the hypotheses of Basic Lemma 2.1, with the same value $k = 1$ in Assumption (3). Hence Lemma 10.1 can be applied to any two of these, and in fact to any finite number taken from this set. Consider the simplest case.

For $A \in \text{Sym}^2(\mathbb{R}^3)$ with eigenvalues $\lambda_1(A) \leq \lambda_2(A) \leq \lambda_3(A)$, and $p(\lambda) \equiv P_1(\lambda_1)P_2(\lambda_2, \lambda_3)$

$$F(A) = P_1(\lambda_1(A))P_2(\lambda_2(A), \lambda_3(A)) = \lambda_1(A) \frac{1}{2}\{\lambda_2(A)^2 + \lambda_3(A)^2\}.$$

The connected component of the set $\{A : F(A) > 0\}$ containing I is $\text{Int } \mathcal{P} \equiv \{A : A > 0\}$. On this set $F > 0$, and $F = 0$ on $\partial \mathcal{P}$. The eigenvalues $\lambda_2(A), \lambda_3(A)$ are positive and $\lambda_2(A)^2 + \lambda_3(A)^2 \geq 2\lambda_1(A)^2$ on this set. Here the determinant majorization is clear.

Appendix A: The interior of the polar cone

Here we describe some useful criteria for determining when a vector is in the interior of the polar of a convex cone. We start with a non-empty open convex cone $\Gamma \subset V$ in a finite dimensional real vector space V . The polar

$$\Gamma^0 \equiv \{y \in V^* : \langle y, x \rangle \geq 0 \ \forall x \in \Gamma\}$$

is a non-empty closed convex cone. As such it has interior when considered as a cone in its vector space span $S \subset V^*$. We denote this relative interior by Γ^* and shall refer to it as the *open polar*, keeping in mind that it is an open convex cone in its span S , which may be have lower dimension than that of V . The orthogonal complement $E \equiv S^\perp$ is called the *edge* of Γ . It is characterized as the linear subspace of Γ which contains all the lines (through the origin) in Γ . Note that if Γ is self-polar (i.e., $\Gamma^0 = \Gamma^0 = \Gamma$), then $\Gamma^* = \Gamma$. In particular, $(\text{Int } \mathcal{P})^* = \text{Int } \mathcal{P}$.

The simple 2-dimensional example $\Gamma \equiv \{x \in \mathbb{R}^2 : x_1 > 0, x_2 > 0\}$, where $E = \{0\}$ and $\Gamma^* = \Gamma$ (self-polar), is a counterexample to the statement:

$$\text{For } y \in S : \quad \langle y, x \rangle > 0 \ \forall x \in \Gamma \quad \implies \quad y \in \Gamma^*,$$

(take $y = (1, 0)$.) However, various strengthenings of this display provide useful criteria for y to belong to the open polar Γ^* .

Lemma A.1 (The Open Polar Criteria). *Suppose $y \in S$. The following are equivalent.*

- (1) $y \in \Gamma^*$.
- (2) $\exists \epsilon > 0$ such that $\langle y, x \rangle \geq \epsilon |x| \quad \forall x \in \overline{\Gamma} \cap S$.
- (3) $\langle y, x \rangle > 0 \quad \forall x \in \overline{\Gamma} - E$.
- (4) $\langle y, x \rangle > 0 \quad \forall x \in (\partial\Gamma - \{0\}) \cap S$.
- (5) $\langle y, x \rangle > 0 \quad \forall x \in \overline{\Gamma} \cap S, x \neq 0$.

Proof (1) \iff (2): Note that:

$$\begin{aligned} y \in \Gamma^* &\iff \text{there exist an } \epsilon - \text{ball } B_\epsilon(y) \subset \Gamma^0 \text{ about } y \\ &\iff \exists \epsilon > 0 \text{ such that } \langle y + \epsilon z, x \rangle \geq 0 \quad \forall |z| \leq 1, x \in \Gamma \\ &\iff \text{(a): } \exists \epsilon > 0 \text{ such that } \langle y, x \rangle \geq \epsilon \langle z, x \rangle \quad \forall |z| \leq 1, x \in \Gamma. \end{aligned}$$

Taking $z = x/|x|$ in (a) yields (2), while (2) yields (a) since $\langle z, x \rangle \leq |x|$ if $|z| \leq 1$.

To see (2) \Rightarrow (3), suppose $x \in \overline{\Gamma} - E$ and decompose x into $x = a + b$ where $a \in E, b \in S$. Then $b \in \overline{\Gamma} \cap S$ and $b \neq 0$, so $\langle y, x \rangle = \langle b, x \rangle > 0$, which proves (3).

That (3) \Rightarrow (4) is obvious.

Since $\overline{\Gamma}$ is convex, it is easy to see that (4) \Rightarrow (5).

To see (5) \Rightarrow (2): We can assume that $E = \{0\}$ and $S = V$. Obviously, (5) \Rightarrow $y \in \Gamma^0$. Hence,

$$\epsilon \equiv \inf_{x \in \overline{\Gamma}, |x|=1} \langle y, x \rangle \geq 0.$$

Since $\overline{\Gamma} \cap \{|x|=1\}$ is compact, there exists $x_0 \in \overline{\Gamma} \cap \{|x|=1\}$ with $\epsilon = \langle y, x_0 \rangle$. Now (5) $\Rightarrow \epsilon > 0$. Hence, $\langle y, \frac{x}{|x|} \rangle \geq \epsilon \quad \forall x \in \overline{\Gamma} - \{0\}$, or $\langle y, x \rangle \geq \epsilon |x| \quad \forall x \in \overline{\Gamma}$. \square

Appendix B: The Gårding gradient map

Let F be a G-D polynomial of degree N on $\text{Sym}^2(\mathbb{R}^n)$, and let Γ be its Gårding cone. We fix $A \in \Gamma$ and write (by Elementary Property (3) in [16])

$$F(A + tB) = F(A) \prod_{j=1}^N (1 + t\lambda_j^{F,A}(B))$$

where $\lambda_j^{F,A}(B)$ are the F -eigenvalues of $B \in \text{Sym}^2(\mathbb{R}^n)$ with respect to $A \in \Gamma$. Hence,

$$\frac{d}{dt} \log F(A + tB) \Big|_{t=0} = \sum_{j=0}^N \lambda_j^{F,A}(B) = \frac{1}{F(A)} \langle (\nabla F)_A, B \rangle \quad (\text{B.1})$$

We shall assume that F is complete which means that all the variables in \mathbb{R}^n are needed to define the G-D operator F . There are many useful equivalent ways of describing completeness (see sect. 3 of [18] for a full discussion of completeness). We point out that an invariant G-D operator is complete.

Here we are using the notation $(\nabla F)_A \equiv D_A F$.

Proposition B.1 *A complete G-D operator F has uniformly elliptic linearization at each point of its Gårding cone Γ . That is, at $A \in \Gamma$*

$$B \mapsto \langle (\nabla F)_A, B \rangle \text{ has coefficient matrix } (\nabla F)_A > 0.$$

Proof We use (3) in Proposition 3.5 in [18] saying that completeness is equivalent to

$$\mathcal{P} \cap E = \{0\}. \quad (\text{B.2})$$

Note that

$$B > 0 \iff \langle B, P \rangle > 0 \ \forall P \in \mathcal{P} - \{0\}. \quad (\text{B.3})$$

Now (B.1) for the derivative of F says that for $A \in \Gamma$ and $P \in \text{Int } \mathcal{P} \subset \Gamma$

$$\langle (\nabla F)_A, P \rangle = F(A) \sum_{j=1}^N \lambda_j^{F,A}(P).$$

Now $F(A) > 0$ and $\lambda_j^{F,A}(P) \geq 0$ for all $P \geq 0$ since $\mathcal{P} \subset \overline{\Gamma}$. Hence, $\langle (\nabla F)_A, P \rangle \geq 0$, and $= 0 \iff \lambda_1^{F,A}(P) = \dots = \lambda_N^{F,A}(P) = 0$, but $\lambda_1^{F,A}(P) = \dots = \lambda_N^{F,A}(P) = 0 \iff P \in E$. Since $P \geq 0$ and $P \in E$ the completeness hypothesis (B.2) implies $P = 0$. Applying (B.3) we have $(\nabla F)_A > 0$. \square

Appendix C: The exhaustion Lemma

Again we take the general point of view. Let $\Gamma \subset V$ be the Gårding cone for a Gårding polynomial \mathbf{g} on a finite dimensional vector space V . It is useful to construct convex *exhaustion functions* $\psi(x)$ for Γ , that is, functions $\psi \in C^\infty(\Gamma)$ which are convex and all the prelevel sets $K_c \equiv \{x \in \Gamma : \psi(x) \leq c\}$ are compact. Now $\psi(x) \equiv -\log \mathbf{g}(x) \in C^\infty(\Gamma)$ is convex and extends to a continuous function on $\overline{\Gamma}$ which is equal to $+\infty$ on $\partial\Gamma$. However, the associated sets $K_c = \{x \in \Gamma : \mathbf{g}(x) \geq e^{-c}\}$ are not compact. For example, when the dimension $n = 1$, if $\mathbf{g}(x) = x$, $\Gamma = (0, \infty) \subset V \equiv \mathbb{R}$, these prelevel sets are not compact. Modifying $-\log x$ by adding the linear function γx with $\gamma > 0$ a constant, yields an exhaustion function ψ for Γ .

There is a caveat here. This scenario given above, of adding a linear function to $\log \mathbf{g}$ does not work if Γ has an edge E . Recall from Appendix A that E is the largest linear subspace contained in Γ . For an extreme example, take the Laplacian $\mathbf{g}(A) = \text{tr}(A)$ on $\text{Sym}^2(\mathbb{R}^n)$ where Γ is the half-space $\{\text{tr}(A) \geq 0\}$ and $E = \{\text{tr}(A) = 0\}$. If we divide by the edge, we fall onto the example above and the argument does work there. In general, the proposed compactness argument holds only after dividing by the edge, or equivalently, restricting to the span $S = E^\perp$. We elect to restrict Γ to the span S since this conforms with the common choice for the complex Monge-Ampère operator, where $S = \text{Sym}^2_{\mathbb{C}}(\mathbb{C}^n)$ is the Hermitian symmetric matrices.

Definition C.1 The convex cone $\Gamma \subset V$ is called regular if the edge of $\overline{\Gamma}$ is $\{0\}$.

The Exhaustion Theorem C.2 Suppose \mathbf{g} is a Gårding polynomial on V of degree N with Gårding cone Γ . Fix $y \in \Gamma^*$, the (relatively) open polar. First, assume Γ is regular. Then the function $\psi(x) \equiv \langle y, x \rangle - \log \mathbf{g}(x)$ is a strictly convex exhaustion function for Γ , so the prelevel sets

$$K_c \equiv \{x \in \Gamma : \langle y, x \rangle - \log \mathbf{g}(x) \leq c\}, \quad \text{for } c \in \mathbb{R}$$

are compact.

If Γ is not regular, then restricting ψ to the span $S \equiv E^\perp$ gives a strictly convex exhaustion function for the regular cone $\Gamma_S \equiv \Gamma \cap S$. Hence, the prelevel sets

$$K_c \equiv \{x \in \Gamma \cap S : \langle y, x \rangle - \log g(x) \leq c\}, \quad \text{for } c \in \mathbb{R}$$

are compact subsets of $\Gamma_S = \Gamma \cap S$.

Proof It suffices to prove the theorem when Γ is regular, since otherwise $g|_S$ has regular Gårding cone $\Gamma_S = \Gamma \cap S$ with the same open polar $\Gamma_S^* = \Gamma^*$.

Recall that $-\log g(x)$ has second derivative at a point $x \in \Gamma$ given by

$$\{D_x^2(-\log g)\}(\xi, \xi) = \sum_{j=1}^N \left(\lambda_j^{g,x}(\xi) \right)^2$$

for all $\xi \in V$, where the Gårding eigenvalues are taken with respect to the direction $x \in \Gamma$. By Gårding [8] (or see [17]), the nullity set

$$\{\xi \in V : \lambda_1^{g,x}(\xi) + \dots + \lambda_N^{g,x}(\xi) = 0\}$$

equals the edge E . Hence, the function $-\log g(x)$ is strictly convex on $S + E^\perp$, and so $\psi(x) \equiv \langle y, x \rangle - \log g(x)$ has the same property since $\langle y, x \rangle$ is affine.

Notice that since $g \equiv 0$ on $\partial\Gamma$ and $\langle y, x \rangle$ is finite, the function $\psi \equiv +\infty$ on $\partial\Gamma$. We conclude that K_c is a closed subset of Γ .

It remains to show that K_c is bounded. For this we use the full hypothesis that y belongs to the open polar Γ^* of Γ , which equals the interior of $\bar{\Gamma}^0$. By Lemma A.1 (2) this is equivalent to the statement

$$(2) \quad \exists \epsilon > 0 \text{ such that } \langle y, x \rangle \geq \epsilon|x| \quad \forall x \in \bar{\Gamma}.$$

This implies that:

$$\begin{aligned} K_c &\subset \{x \in \bar{\Gamma} : e^{\epsilon|x|} \leq e^c g(x)\} \subset \left\{x \in \bar{\Gamma} : \frac{(\epsilon|x|)^{N+1}}{(N+1)!} \leq e^c g(x)\right\} \\ &= \left\{x \in \bar{\Gamma} : |x| \leq \frac{(N+1)!e^c}{\epsilon^{N+1}} g\left(\frac{x}{|x|}\right)\right\} \subset B_R(0) \end{aligned}$$

with

$$R \equiv \frac{(N+1)!e^c}{\epsilon^{N+1}} \sup_{\xi \in \bar{\Gamma}, |\xi|=1} g(\xi) < \infty.$$

□

Corollary C.3 For constants $c_1, c_2 \in \mathbb{R}$ and $y \in \Gamma^*$, the set

$$K_{c_1, c_2} \equiv \{x \in \Gamma : \langle y, x \rangle \leq c_1 \text{ and } \log g(x) \geq c_2\}$$

is compact.

Proof One has $K_{c_1, c_2} \subset K_{c_1 - c_2}$. □

Appendix D: The Güler derivative estimates for G-D operators

Suppose F is a Gårding-Dirichlet operator on $\text{Sym}^2(\mathbb{R}^n)$ of degree N with Gårding cone Γ . (We assume $F \not\equiv 0$, and recall that by definition Γ is open.) For each $A \in \Gamma$ there is a simple formula for the k^{th} derivative of $L \equiv \log F$ at A in terms of the Gårding eigenvalues $\lambda_j^{F,A}$ of F w.r.t. the direction A .

In this section we shall abbreviate $\lambda_j^{F,A}$ to λ_j^A or just λ_j .

Theorem D.1 (Derivatives). *At a point $A \in \Gamma$ and $\forall B \in \text{Sym}^2(\mathbb{R}^n)$:*

(1)

$$\langle D_A F, B \rangle = F(A) \sum_{j=1}^N \lambda_j^A(B) \quad \text{and}$$

$$\langle D_A F, B \rangle \geq 0 \quad \forall B \in \overline{\Gamma}, \quad \text{with equality} \iff B \in E.$$

(2)

$$D_A^2(\log F)(B, B) = - \sum_{j=1}^N (\lambda_j^A(B))^2 \quad \text{and}$$

$$\text{the quadratic form } D_A^2(\log F) \leq 0 \quad \text{with null space } E.$$

(3)

$$D_A^3(\log F)(B, B, B) = 2 \sum_{j=1}^N (\lambda_j^A(B))^3$$

(4)

$$D_A^{(k)}(\log F)(B, \dots, B) = (-1)^{k-1} (k-1)! \sum_{j=1}^N (\lambda_j^A(B))^k \quad \forall k \geq 1$$

In particular,

(1)' The directional derivative of F at A is strictly increasing in all directions B in the closed Gårding cone $\overline{\Gamma}$, other than the edge directions where it is zero.

(2)' The restriction of $L = \log F$ to the span S is a strictly concave operator on $\Gamma_S \equiv S \cap \Gamma$.

Proof For $A \in \Gamma$ and $B \in \text{Sym}^2(\mathbb{R}^n)$, let $\varphi(t) \equiv \log F(A + tB)$. Then

$$F(A + tB) = F(A) \prod_{j=1}^N (1 + t\lambda_j^A(B)).$$

(See for example [8] or the Elementary Property (3) in [16].)

Since $\varphi(t) = \log F(A) + \sum_{j=1}^N \log(1 + t\lambda_j^A)$, we have (with $\lambda_j \equiv \lambda_j^A$) that

$$\varphi'(t) = \sum_{j=1}^N \frac{\lambda_j}{1 + t\lambda_j}, \quad \varphi''(t) = - \sum_{j=1}^N \frac{\lambda_j^2}{(1 + t\lambda_j)^2},$$

and for $k \geq 2$,

$$\varphi^{(k)}(t) = (-1)^{k-1}(k-1)! \sum_{j=1}^N \frac{\lambda_j^k}{(1+t\lambda_j)^k}.$$

Since $\varphi^{(k)}(0) = \{D_A^{(k)}(\log F)\}(B, \dots, B)$ this proves all the equalities in (1), (2), (3) and (4). The remainder of the theorem follows from the fact that for all $B \in \overline{\Gamma}$, the eigenvalues $\lambda_1^A(B), \dots, \lambda_N^A(B)$ are all ≥ 0 with equality iff $B \in E$, that is, the nullity $\{B : \lambda_1^A(B) = \dots = \lambda_N^A(B) = 0\}$ equals the edge E (see [8, 16]).

Now with $\lambda_j \equiv \lambda_j^A(B) \equiv \lambda_j^{F,A}(B)$ and $\lambda \equiv (\lambda_1, \dots, \lambda_N) \in \mathbb{R}^N$, the little ℓ^k -norm of λ is

$$\|\lambda\|_k \equiv \left(\sum_{j=1}^N \|\lambda_j\|^k \right)^{\frac{1}{k}} \quad \text{for } k \geq 1 \quad \text{and} \quad \|\lambda\|_\infty = \sup_{1 \leq j \leq N} |\lambda_j|.$$

Recall that

$$\|\lambda\|_k \leq \|\lambda\|_\ell \quad \text{for } \ell = 1, 2, \dots, k-1 \text{ with equality iff } \lambda \text{ is an axis vector.} \quad (\text{D.1})$$

To see this we can assume that $\lambda \neq 0$. By homogeneity one can assume $\|\lambda\|_\ell = 1$, and therefore $0 \leq |\lambda_j| \leq 1$ for all j . Then $\ell < k$ implies $\sum |\lambda_j|^k \leq \sum |\lambda_j|^\ell = 1$. Hence, $\|\lambda\|_k \leq 1 = \|\lambda\|_\ell$ with equality if and only if λ is a unit axis vector.

This translates into an upper bound estimate for the k^{th} derivative by lower order derivatives.

Theorem D.2 (Güler [9]). *For $A \in \Gamma$, $B \in \text{Sym}^2(\mathbb{R}^n)$ and k fixed, if $\ell = 2, 4, \dots$ is even and $\ell \leq k$, then*

$$\left| \frac{1}{(k-1)!} \left\{ D_A^{(k)} \log F \right\} (B, \dots, B) \right|^{\frac{1}{k}} \leq \left[\frac{1}{(\ell-1)!} \left\{ D_A^{(\ell)} \log F \right\} (B, \dots, B) \right]^{\frac{1}{\ell}}$$

Proof Note that for $k = 1, 3, \dots$ odd,

$$\left| \frac{1}{(k-1)!} \left\{ D_A^{(k)} \log F \right\} (B, \dots, B) \right| = \left| \sum_{j=1}^N \lambda_j^k \right| \leq \sum_{j=1}^N |\lambda_j|^k = \|\lambda\|_k^k,$$

and for $k = 2, 4, \dots$ even

$$\frac{1}{(k-1)!} \left\{ D_A^{(k)} \log F \right\} (B, \dots, B) = \|\lambda\|_k^k.$$

Hence, in both cases, $\|\lambda\|_k \leq \|\lambda\|_\ell$ for $\ell = 1, \dots, k-1$ yields the estimates for the k^{th} derivative of $\log F$. \square

For equality to hold in Theorem D.2, B must either be in the edge E or have exactly one non-zero A -eigenvalue.

Further discussion–second derivative formulas

The above first and second derivative formulas for $\log F$ at $A \in \Gamma$, namely

$$(1) \quad \langle D_A \log F, B \rangle = \sum_{j=1}^N \lambda_j \quad \text{and} \quad (2) \quad (D_A^2 \log F)(B, B) = - \sum_{j=1}^N \lambda_j^2$$

for all $B \in \text{Sym}^2(\mathbb{R}^n)$ can be used to conclude interesting formulas for the second derivatives of F and $F^{\frac{1}{N}}$. Here we abbreviate $\lambda_j^{F,A}(B)$ to just λ_j or $\lambda_j(B)$, and we recall the discriminant polynomial

$$\text{Discr}(\lambda_1, \dots, \lambda_N) = \sum_{i < j} (\lambda_i - \lambda_j)^2.$$

Proposition D.3

- (D.1) $(D_A^2 \log F)(B, B) = -|\lambda(B)|^2$
- (D.2) $(D_A^2 F)(B, B) = 2F(A)\sigma_2(\lambda(B))$
- (D.3) $(D_A F^{\frac{1}{N}})(B, B) = -\frac{1}{N^2} F(A)^{\frac{1}{N}} \text{Discr}(\lambda(B))$

Moreover, the quadratic forms $D_A^2 \log F$ and $D_A^2 F$ have the same null space, namely $N \equiv \{B : \lambda_1(B) = \dots = \lambda_N(B) = 0\}$. The nullity N is the same as the edge $E \equiv \bar{\Gamma} \cap (-\bar{\Gamma})$ and also the linearity of F which is, by definition, the largest linear subspace on which $F = 0$. Modulo this subspace, $D_A^2 \log F < 0$, while $D_A^2 F$ has Lorentzian signature with future light cone Γ . Finally, the quadratic form $\text{Discr}(\lambda(B)) \geq 0$ with null space $\{B \in \text{Sym}^2(\mathbb{R}^n) : \lambda_1(B) = \dots = \lambda_N(B)\} = E + \mathbb{R} \cdot A$ of dimension $= \dim(E) + 1$.

Note that (D.3) proves the Gårding Lemma that $F^{\frac{1}{N}}$ is concave.

Proof of (D.2) By (1) and (2) above,

$$\begin{aligned} - \sum_{j=1}^N \lambda_j^2 &= D_A^2 \log F = D_A \left(\frac{D_A F}{F(A)} \right) = \frac{D_A^2(F)}{F(A)} \\ &= - \frac{D_A F \circ D_A F}{F(A)^2} = \frac{D_A^2 F}{F(A)} - \left(\sum_{j=1}^N \lambda_j \right)^2. \end{aligned}$$

Note that $(\sum_j \lambda_j)^2 - \sum_j \lambda_j^2 = 2\sigma_2(\lambda)$. □

Sketch of proof of (D.3) Use the standard formula (obtained by expanding out the right hand term)

$$N|\lambda(B)|^2 - \sigma_1(\lambda(B))^2 = \text{Discr}(\lambda(B)) \equiv \sum_{i < j} (\lambda_i(B) - \lambda_j(B))^2$$

to compute that

$$(D_A^2 F^{\frac{1}{N}})(B, B) = -\frac{1}{N^2} F^{\frac{1}{N}}(A) \left\{ \text{Discr}(\lambda^A(B)) \right\}.$$

Appendix E: The central ray hypothesis

Suppose F is a G-D operator on $\text{Sym}^2(\mathbb{R}^n)$ of degree N with Gårding cone Γ . (We assume $F \not\equiv 0$ and by definition Γ is open and contains $\{A : A > 0\}$.)

The next lemma shows that there exists a unique ray \mathcal{R} contained in the Gårding cone Γ , with the property that Γ is, in a certain sense, symmetrical about \mathcal{R} .

Lemma E.1

$$F(A)^{\frac{1}{N}} \leq \left[\sup_{B \in \overline{\Gamma}, \|B\|=1} F(B)^{\frac{1}{N}} \right] \|A\| \quad \text{for all } A \in \overline{\Gamma} - E. \quad (\text{E.1})$$

This sup is attained at a unique maximum point B_0 which belongs to $\Gamma_S \equiv \Gamma \cap S$ where S is the span. Equivalently, equality occurs at a point $A \in \overline{\Gamma} \iff A \in \mathcal{R} \equiv \mathbb{R}^+ \cdot B_0$, the ray through B_0 .

Definition E.2 This unique ray \mathcal{R} for (F, Γ) is called the *central ray*.

Proof Suppose B_0 is a maximum point. First, $B_0 \in \Gamma$ since F vanishes on $\partial\Gamma$. Secondly, $B_0 \in S$. If not, then writing $B_0 = B_E + B_S$ with respect to the orthogonal decomposition $\text{Sym}^2(\mathbb{R}^n) = E \oplus S$, we have $B_E \neq 0$. Therefore, $\|B_S\| < \|B_0\| = 1$, and hence

$$F\left(\frac{B_S}{\|B_S\|}\right)^{\frac{1}{N}} = \frac{1}{\|B_S\|} F(B_S)^{\frac{1}{N}} = \frac{1}{\|B_S\|} F(B_0)^{\frac{1}{N}} > F(B_0)^{\frac{1}{N}}.$$

which contradicts that B_0 is a maximum point.

For the remainder of the proof, we can restrict to the span $S = E^\perp$, and we set $\gamma \equiv \sup_{B \in \overline{\Gamma}, \|B\|=1} F(B)^{\frac{1}{N}}$.

The tangent affine hyperplane $H \equiv \{A : \langle B_0, A \rangle = 1\}$ separates the level sets $\{F(B)^{\frac{1}{N}} = \gamma\}$ and $\{\|B\| = 1\}$ since $F^{\frac{1}{N}}$ is strictly concave, and $\|B\|$ is strictly convex on S . Furthermore, B_0 is the only point of intersection of these hypersurfaces, since H touches each of these hypersurfaces only at the point B_0 .

If the sup in (E.1) is attained at a point A we may assume $\|A\| = 1$ by dividing the inequality by $\|A\|$. Now we must have $A = B_0$ since all other points on the unit sphere are strictly below the hypersurface $F^{\frac{1}{N}} = \gamma$. \square

Corollary E.3 There is a unique point A_0 , up to positive multiples, where the sphere and the level set of $F^{\frac{1}{N}}$ have a common normal.

Proof Let A_0 be a point where there is a common normal. We can renormalize so that $\|A_0\| = 1$, since everything is of degree 1. Let H_0 be the tangent hyperplane to the sphere at A_0 . Then as in the proof of the theorem above H_0 strictly separates the sphere and the hypersurface $F(A)^{\frac{1}{N}} = F(A_0)^{\frac{1}{N}}$ away from A_0 ; and the value of $F^{\frac{1}{N}}$ at all points $\neq A_0$ in the sphere are $< F(A_0)^{\frac{1}{N}}$. Thus if $A_0 \neq B_0$ we would have $F(A_0)^{\frac{1}{N}} > F(B_0)^{\frac{1}{N}} > F(A_0)^{\frac{1}{N}}$. \square

Corollary E.4 The central ray \mathcal{R} is characterized by the fact that for any non-zero point $B \in \mathcal{R}$, $D_B F$ and $D_B \|A\|^2 = 2B$ are positively proportional, i.e.,

$$D_B F = kB$$

with $B \in \Gamma_S = S \cap \Gamma$ and $k > 0$.

We now examine conditions which imply that the central ray is $\mathbb{R}_+ I$.

Proposition E.5 *If F is invariant under $O(n)$, $SU(m)$ with $n = 2m$, Sp_m with $n = 4m$, or any other subgroup of $O(n)$ whose only fixed line in $\text{Sym}^2(\mathbb{R}^n)$ is the one through the identity I , then the central ray \mathcal{R} is the ray through I .*

Proof If F is invariant under a subgroup $G \subset O(n)$, then G fixes the central ray. \square

Definition E.6 The *Central Ray Hypothesis* (CRH) is that: The identity I generates the central ray \mathcal{R} of (Γ, F) .

There are several important ways of formulating the CRH, which we now describe.

Definition E.7 The *Gårding (or F -) Laplacian*, denoted Δ^F , is defined by

$$\Delta^F(B) \equiv \sum_{j=1}^N \lambda_j^{F,I}(B) \equiv \text{tr}^F(B).$$

Note that, after normalizing $F(I) = 1$,

$$\begin{aligned} \Delta^F(B) &= \frac{d}{dt} \Big|_{t=0} F(I + tB) = \frac{1}{(N-1)!} \frac{d^{(N-1)}}{dt^{(N-1)}} \Big|_{t=0} F(tI + B) \\ &= \sigma_1^F(B) = \langle D_I \log F, B \rangle. \end{aligned}$$

Theorem E.8 (Equivalent Formulations of the CRH). *The following conditions are equivalent.*

(1)

$$F(A)^{\frac{1}{N}} \leq \left[\sup_{\|B\|^2=n, B \in \Gamma} F(B)^{\frac{1}{N}} \right] \|A\| \quad \forall A \in \Gamma$$

with equality iff $A = tI, t > 0$.

(2) $D_I \log F = \frac{1}{F(I)} D_I F = kI$, for $k > 0$.

(3) $\Delta^F = k\Delta^{\text{standard}}$, for $k > 0$.

(4) $F(A)^{\frac{1}{N}} \leq F(I)^{\frac{1}{N}} \frac{k}{N} \langle I, A \rangle \quad \forall A \in \Gamma \quad \text{with equality iff } A \in tI + E$

for $t > 0$. The k, n, N are related by $kn = N$.

Proof We have already shown that (1) \iff (2). Now (2) \iff (3) since $\Delta^F(B) = \langle D_I \log F, B \rangle$ implies that $\Delta^F = k\Delta^{\text{standard}} \iff D_I \log F = kI$. Statement (4) says that $F^{\frac{1}{N}}(A)$ is bounded above by $c\langle I, A \rangle$ with equality at $tI + E, t > 0$. Thus the derivative of $F^{\frac{1}{N}}$ at I is a multiple of I which is equivalent to (3). \square

References

1. Abja, S., Olive, G.: Local regularity for concave homogeneous complex degenerate elliptic equations dominating the Monge-Ampère equation. *Ann. Mat. Pura Appl.* **201**(2), 561–587 (2022)
2. Abja, S., Dinev, S., Olive, G.: Uniform estimates for concave homogeneous complex degenerate elliptic equations comparable to the Monge-Ampère equation. *Potential Anal.* (2022). <https://doi.org/10.1007/s11118-022-10009-w>

3. Alesker, S., Verbitsky, M.: Plurisubharmonic functions on hypercomplex manifolds and HKT-geometry, [arXiv:0510140v3](https://arxiv.org/abs/0510140v3)
4. Bauschke, H.H., Güler, O., Lewis, A.S., Sendov, H.S.: Hyperbolic polynomials and convex analysis. *Canad. J. Math.* **53**(3), 470–488 (2001)
5. Caffarelli, L., Cabré, X.: Fully nonlinear elliptic equations. Colloquium Publications, American (1995)
6. Caffarelli, L., Nirenberg, L., Spruck, J.: The Dirichlet problem for nonlinear second-order elliptic equations. III. Functions of the eigenvalues of the Hessian. *Acta Math.* **155**(3–4), 261–301 (1985)
7. Dinew, S.: Interior estimates for p -plurisubharmonic functions, [arXiv:2006.12979](https://arxiv.org/abs/2006.12979)
8. Gårding, L.: An inequality for hyperbolic polynomials. *J. Math. Mech.* **8**, 957–965 (1959)
9. Güler, O.: Hyperbolic polynomials and interior point methods for convex programming. *Math. Oper. Res.* **22**(2), 350–377 (1997)
10. Guo, B., Phong, D.H., Tong, F.: On L^∞ -estimates for complex Monge-Ampère equations, [arXiv:2106.02224](https://arxiv.org/abs/2106.02224)
11. Guo, B., Phong, D. H.: On L^∞ -estimates for fully nonlinear partial differential equations on hermitian manifolds, [arXiv:2204.12549v1](https://arxiv.org/abs/2204.12549v1)
12. Guo, B., Phong, D. H.: Uniform entropy and energy bounds for fully non-linear equations, (2022) [arXiv:2207.08983v1](https://arxiv.org/abs/2207.08983v1)
13. Gurvits, L.: Van der Waerden/Schrijver-Valiant like conjectures and stable (aka hyperbolic) homogeneous polynomials: one theorem for all. *Electron. J. Comb.* **15**(1), 26 (2008)
14. Harvey, F.R., Lawson, H.B.: Dirichlet duality and the nonlinear Dirichlet problem. *Comm. Pure Appl. Math.* **62**(3), 396–443 (2009)
15. Harvey, F.R., Lawson, H.B.: Dirichlet duality and the nonlinear Dirichlet problem on Riemannian manifolds. *J. Diff. Geom.* **88**(3), 395–482 (2011)
16. Harvey, F.R., Lawson, H.B.: Hyperbolic polynomials and the Dirichlet problem, [ArXiv:0912.5220](https://arxiv.org/abs/0912.5220)
17. Harvey, F.R., Lawson, H.B.: Gårding's theory of hyperbolic polynomials. *Comm. Pure App. Math.* **66**(7), 1102–1128 (2013)
18. Harvey, F.R., Lawson, H.B.: Lagrangian potential theory and a Lagrangian equation of Monge-Ampère type. In: Cao, H.-D., Li, J., Schoen, R., Yau, S.-T. (eds.) *Surveys in Differential Geometry*, pp. 217–257. International Press, Somerville (2018)
19. Harvey, F.R., Lawson, H.B.: The inhomogeneous Dirichlet problem for natural operators on manifolds. *Ann. l'Inst. Fourier* **69**(7), 3017–3064 (2019). [arXiv:1805.11121](https://arxiv.org/abs/1805.11121)
20. Wang, Y.: On the $C^{2,\alpha}$ -regularity of the complex Monge-Ampère equation. *Math. Res. Lett.* **19**(4), 939–946 (2012)
21. Yau, S.T.: On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation. I. *Comm. Pure Appl. Math.* **31**, 339–411 (1978)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.